JAWAID AKHTAR, I.A.S.,

Principal Secretary to Government Health and Family Welfare Department and Medical Education Department



Tel: 080-2225 5324 080-2203 4234

Fax: 080-2235 3916

E-mail: prs-hfw@karnataka.gov.in Room No. 105, First Floor Vikasa Soudha, Dr. B.R. Ambedkar Veedhi

Bengaluru - 560 001

Date: 7.04.2020

No: HFW 89 ACS 2020

ORDER

Sub: Revised Lab Testing Protocol- Karnataka (7th April 2020)

Orders have been issued from time to time by Additional Chief Secretary-Health and Family welfare and Commissioner-Health and Family welfare services regarding lab testing protocols for COVID-19. In view of recent revision of protocol by ICMR-GOI and superseding all the previous circulars, below revised guidelines are being issued.

A. Indications for RT-PCR (Throat swab + Nasal Swab):

- I. All symptomatic individuals who have undertaken international travel in the last 14 days
- II. All primary contacts (direct and high-risk contacts) of Confirmed cases
 - a. If symptomatic Test immediately
 - b. If Asymptomatic -should be **tested once** between day 5 and day 14 of coming in his/her contact
- III. All symptomatic health care workers
- IV. All hospitalized patients with Severe Acute Respiratory Illness (fever AND cough and/or shortness of breath)
- V. ILI patients in containment zone (Once Rapid Antibody test commences, such patients will be tested through rapid kits)
- VI.ILI Patients in whom Rapid test is negative and clinician warrants RT-PCR, test can be done

The kit for Rapid Antibody tests will be received shortly and tests will commence. The protocol for Rapid Antibody tests will be as follows:-

B. Indications for Rapid Antibody based Blood Test:

Symptomatic Influenza Like illness-ILI

(Cough, Cold, low grade fever, sore throat)

- I. Patients coming to designated Fever clinics
- II. Patients identified during house-to-house survey in containment zone

III. Patients in large migratory gatherings/shelters

IV. Above patients in whom first Rapid test has come negative and RT-PCR not done, Repeat Rapid test 10 days after first Rapid test.

General Guidelines:

- Healthcare workers doing the rapid antibody test to use gloves, N-95 mask, and headcovers.
- Health care workers collecting throat/nasal swab to use complete PPE and should collect samples at designated facility
- The rapid antibody tests approved by US-FDA/CE-IVD or non-CE-IVD validated by ICMR-NIV with marketing approval by DCGI only to beused.
- In order to ensure that all such cases are monitored and necessary action is initiated with respect to infectious disease management, details of all test results shall be uploaded inICMR portal.
- As a matter of abundant precautions, all symptomatic ILI persons should be advised home quarantine for 14 days
- When home quarantine is not feasible, consider facility-based quarantine

(JAWAID AKHTAR)

Additional Chief Secretary to Government
Health and Family Welfare Department

- 1. Commissioner, Health & Family Welfare Services, Anand Rao Circle, Bengaluru.
- 2. Mission Director, National Health Mission, Anand Rao Circle, Bengaluru.
- 3. Director, Health & Family Welfare Services, Anand Rao Circle, Bengaluru.
- 4. Director, Medical Education Dept., Anand Rao Circle, Bengaluru.
- 5. All Deputy Commissioners.
- 6. All CEOs of ZPs.
- 7. All District Health Officers.
- 8. All District Surgeons.
- 9. All Testing Labs of State.

Copy for kind information:

Chief Secretary, Government of Karnataka.